

# LIST OF REFERENCES CITED BY APPLICANT

(Use several sheets if necessary)

ATTY. DOCKET NO.

7853-238

APPLICATION NO.

TBA

APPLICANT


Kingsbury et al.

FILING DATE

Herewith

GROUP

1644



## U.S. PATENT DOCUMENTS

EXAMINER INITIAL		DOCUMENT NUMBER	DATE	NAME	CLASS	SUBCLASS	FILING DATE IF APPROPRIATE
ay	AA	5,840,691	11/24/98	Furcht et al.			
ay	AB	5,721,351	2/24/98	Levinson			
ay	AC	5,591,719	1/7/97	Furcht et al.			
ay	AD	5,262,311	11/16/93	Pardee and Liang			
ay	AE	5,116,964	5/26/92	Capon and Lasky			

## FOREIGN PATENT DOCUMENTS

		DOCUMENT NUMBER	DATE	COUNTRY	CLASS	SUBCLASS	TRANSLATION	
							YES	NO
	AF	WO 99/15663	4/1/99	PCT				
	AG	WO 97/42224	11/13/97	PCT				
	AH	JP 6-178687	6/28/94	Japan				
	AI	EP 239400	8/3/94	EPO				




## OTHER REFERENCES (Including Author, Title, Date, Pertinent Pages, Etc.)

	AJ	GenBank Accession No. E07714 (cDNA encoding mouse ST2 protein)
	AK	GenBank Accession No. P14719 (mouse ST2 protein precursor)
	AL	GenBank Accession No. E08652 (cDNA encoding mouse ST2L)
	AM	GenBank Accession No. S29498 (mouse ST2L protein)
	AN	GenBank Accession No. AB012701 (human mRNA for ST2L, 5'noncoding region incl. exon 1a, complete cds)
	AO	GenBank Accession No. BAA82405 (human ST2L protein)
	AP	GenBank Accession No. D12763 (human mRNA for ST2L protein)
	AQ	GenBank Accession No. BAA0223 (human ST2 protein)
	AR	GenBank Accession No. AB029084 (human ST2V mRNA, complete cds)
	AS	GenBank Accession No. BAA85894 (human ST2V)
	AT	GenBank Accession No. Y07519 (mouse St2 gene)
	AU	GenBank Accession No. M28621 (mouse interferon-γ mRNA sequence)
	AV	Altschul et al., 1997, "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs", Nucl Acids Res 25:3389-3402
	AW	Altschul et al., 1990, "Basic Local Alignment Search Tool", J. Mol. Biol. 215:403-410
	AX	Anderson and Coyle, 1994, "TH2 and 'TH2-like' cells in allergy and asthma: pharmacological perspectives", Trends Pharmacol. Sci. 15:324-332

AY	Aruffo et al., 1990, "CD44 is the principal cell surface receptor for hyaluronate", <i>Cell</i> <b>61</b> :1303-1313
AZ	Bergers et al., 1994, "Alternative promoter usage of the Fos-responsive gene <i>Fit-1</i> generates mRNA isoforms coding for either secreted or membrane-bound proteins related to the IL-1 receptor", <i>EMBO J.</i> <b>13</b> :1176-1188
BA	Chen et al., 1993, "RAG-2-deficient blastocyst complementation: an assay of gene function in lymphocyte development", <i>Proc. Natl. Acad. Sci. USA</i> <b>90</b> :4528-4532
BB	Clerici et al., 1993, "Restoration of HIV-specific cell-mediated immune responses by interleukin-12 in vitro", <i>Science</i> <b>262</b> :1721-1724
BC	Clerici et al., 1993, "Changes in interleukin-2 and interleukin-4 production in asymptomatic, human immunodeficiency virus-seropositive individuals", <i>J. Clin. Invest.</i> <b>91</b> :759-765
BD	Cohn et al., 1997, "Induction of airway mucus production by T helper 2 (Th2) cells: a critical role for interleukin 4 in cell recruitment but not mucus production", <i>J. Exp. Med.</i> <b>186</b> :1737-1747
BE	Del Prete et al., 1991, "Purified protein derivative of <i>Mycobacterium tuberculosis</i> and excretory-secretory antigen(s) of <i>Toxocara canis</i> expand in vitro human T cells with stable and opposite (Type 1 T helper or Type 2 T helper) profile of cytokine production", <i>J. Clin. Invest.</i> <b>88</b> :346-350
BF	Fägerstam et al., 1992, "Biospecific interaction analysis using surface plasmon resonance detection applied to kinetic, binding site and concentration analysis", <i>J. Chromatog.</i> <b>597</b> :397-410
BG	Firestein et al., 1989, "A new murine CD4 <sup>+</sup> T cell subset with an unrestricted cytokine profile", <i>J. Immunol.</i> <b>143</b> :518-525
BH	Frangogiannis et al., 1998, "Resident cardiac mast cells degranulate and release preformed TNF- $\alpha$ initiating the cytokine cascade in experimental canine myocardial ischemia/reperfusion", <i>Circulation</i> <b>98</b> :699-710
BI	Gächter et al., 1996, "Transcription of the interleukin-1 receptor-related T1 gene is initiated at different promoters in mast cells and fibroblasts", <i>J. Biol. Chem.</i> <b>271</b> :124-129
BJ	Gavett et al., 1994, "Depletion of murine CD4 <sup>+</sup> T lymphocytes prevents antigen-induced airway hyperreactivity and pulmonary eosinophilia", <i>Am. J. Respir Cell Mol. Biol.</i> <b>10</b> :587-593
BK	George et al., 1994, "Embryonic expression and cloning of the murine GATA-3 gene", <i>Development</i> <b>120</b> :2673-2686
BL	Gonazlo et al., 1996, "Mouse eotaxin expression parallels eosinophil accumulation during lung allergic inflammation but it is not restricted to a Th2-type response", <i>Immunity</i> <b>4</b> :1-14
BM	Gu et al., 1994, "Deletion of a DNA polymerase $\beta$ gene segment in T cells using cell type-specific gene targeting", <i>Science</i> <b>265</b> :103-106
BN	Hamelmann et al., 1997, "Noninvasive measurement of airway responsiveness in allergic mice using barometric plethysmography", <i>Am. J. Respir. Crit. Care Med.</i> <b>156</b> :766-775
BO	Holgate, 1997, "The cellular and mediator basis of asthma in relation to natural history", <i>Lancet</i> <b>350</b> (Suppl. 2):5-9
BP	Kanagawa et al., 1993, "Resistance of mice deficient in IL-4 to retrovirus-induced immunodeficiency syndrome (MAIDS)", <i>Science</i> <b>262</b> :240-242
BQ	Kaneshima et al., 1994, "Human hematolymphoid cells in SCID mice", <i>Curr. Opin. Immunol.</i> <b>6</b> :327-333

BR	Karlin and Altschul, 1993, "Applications and statistics for multiple high-scoring segments in molecular sequences", Proc. Natl. Acad. Sci. USA <u>90</u> :5873-5877
BS	Karlin and Altschul, 1990, "Methods for assessing the statistical significance of molecular sequence features by using general scoring schemes", Proc Natl Acad Sci USA <u>87</u> :2264-2268
BT	Klemenz et al., 1989, "Serum- and oncoprotein-mediated induction of a gene with sequence similarity to the gene encoding carcinoembryonic antigen", Proc. Natl. Acad. Sci. USA <u>86</u> :5708-5712
BU	Ko and Engel, 1993, "DNA binding specificities of the GATA transcription factor family", Mol. Cell. Biol. <u>13</u> :4011-4022
BV	Kumar et al., 1997, "Expression of ST2, an interleukin-1 receptor homologue, is induced by proinflammatory stimuli", Biochem. Biophys. Res. Comm. <u>235</u> :474-478
BW	Liew, 1994, "Induction and regulation of CD4 <sup>+</sup> T cell subsets", CIBA Foundation Symposium <u>87</u> :170-178
BX	Löhning et al., 1998, "T1/ST2 is preferentially expressed on murine Th2 cells, independent of interleukin 4, interleukin 5, and interleukin 10, and important for Th2 effector function", Proc. Natl. Acad. Sci. USA <u>95</u> :6930-6935
BY	Lukacs et al., 1994, "Interleukin-4-dependent pulmonary eosinophil infiltration in a murine model of asthma", Am. J. Respir. Cell Mol. Biol. <u>10</u> :526-532
BZ	Maggi et al., 1994, "Th2-like CD8 <sup>+</sup> T cells showing B cell helper function and reduced cytolytic activity in human immunodeficiency virus type 1 infection", J. Exp. Med. <u>180</u> :489-495
CA	Maggi et al., 1994, "Ability of HIV to promote a T <sub>H</sub> 1 to T <sub>H</sub> 0 shift and to replicate preferentially in T <sub>H</sub> 2 to T <sub>H</sub> 0 cells", Science <u>265</u> :244-248
CB	Makino et al., 1990, "H-2-associated and background genes influence the development of a murine retrovirus-induced immunodeficiency syndrome", J. Immunol. <u>144</u> :4347-4355
CC	Manetti et al., 1994, "CD30 expression by CD8 <sup>+</sup> T cells producing type 2 helper cytokines. Evidence for large numbers of CD8 <sup>+</sup> CD30 <sup>+</sup> T cell clones in human immunodeficiency virus infection", J. Exp. Med. <u>180</u> :2407-2411
CD	McMahan et al., 1991, "A novel IL-1 receptor, cloned from B cells by mammalian expression, is expressed in many cell types", EMBO J. <u>10</u> :2821-2832
CE	Metzler and Xu, 1997, "The role of mast cells in atherosclerosis", Int. Arch. Allergy Immunol. <u>114</u> :10-14
CF	Moll et al., 1991, "Expression of T-cell-associated serine proteinase 1 during murine <i>Leishmania major</i> infection correlates with susceptibility to disease", Infect. Immun. <u>59</u> :4701-4705
CG	Mosmann and Moore, 1991, "The role of IL-10 in crossregulation of T <sub>H</sub> 1 and T <sub>H</sub> 2 responses", Immunol Today <u>12</u> :A49-A53
CH	Mosmann and Coffman, 1989, "TH1 and TH2 cells: Different patterns of lymphokine secretion lead to different functional properties", Annu. Rev. Immunol. <u>7</u> :145-173
CI	Murphy et al., 1990, "Induction by antigen of intrathymic apoptosis of CD4 <sup>+</sup> CD8 <sup>+</sup> TCR <sup>+</sup> thymocytes in vivo", Science <u>250</u> :1720-1723
CJ	Myers and Miller, 1988, "Optimal Alignments in Linear Space", Comput. Appl. Biosci. <u>4</u> :11-17

CK	Oettgen and Gheea, 1999, "IgE in asthma and atopy: cellular and molecular connections", J. Clin. Invest. <u>104</u> :829-835
CL	Onashi et al., 1992, "Airway hyperresponsiveness, increased intracellular spaces of bronchial epithelium, and increased infiltration of eosinophils and lymphocytes in bronchial mucosa in asthma", Am. Rev Respir. Dis. <u>145</u> :1469-1476
CM	Ray and Cohn, 1999, "Th2 cells and GATA-3 in asthma: new insights into the regulation of airway inflammation", J. Clin. Invest. <u>104</u> :985-993
CN	Robinson et al., 1993, "Activation of CD4 <sup>+</sup> T cells, increased T <sub>H2</sub> -type cytokine mRNA expression, eosinophil recruitment in bronchoalveolar lavage after allergen inhalation challenge in patients with atopic asthma", J. Allergy Clin. Immunol. <u>92</u> :313-324
CO	Romagnani, 1997, "The Th1/Th2 paradigm", Immunol. Today <u>18</u> :263-266
CP	Schweitzer and Sharpe, 1998, "Studies using antigen-presenting cells lacking expression of both B7-1 (CD80) and B7-2 (CD86) show distinct requirements for B7 molecules during priming versus restimulation of Th2 but not Th1 cytokine production", J. Immunol. <u>161</u> :2762-2771
CQ	Seder et al., 1994, "CD28-mediated costimulation of interleukin 2 (IL-2) production plays a critical role in T cell priming for IL-4 and interferon- $\gamma$ production", J. Exp. Med. <u>179</u> :299-304
CR	Seder and Gros, 1995, "The functional role of CD8 <sup>+</sup> T helper type 2 cells", J. Exp. Med. <u>181</u> :5-7
CS	Shinkai et al., 1992, "RAG-2-deficient mice lack mature lymphocytes owing to inability to initiate V(D)J rearrangement", Cell <u>68</u> :855-867
CT	Tepper et al., 1990, "IL-4 induces allergic-like inflammatory disease and alters T cell development in transgenic mice", Cell <u>62</u> :457-467
CU	Tominaga et al., 1999, "Presence and expression of a novel variant form of ST2 gene product in human leukemic cell line UT-7/GM", Biochem. Biophys. Res. Comm. <u>264</u> :14-18
CV	Tominaga et al., 1992, "Nucleotide sequence of a complementary DNA for human ST2", Biochim. Biophys. Acta <u>1171</u> :215-218
CW	Tominaga et al., 1991, "Molecular cloning of the murine ST2 gene. Characterization and chromosomal mapping", Biochim. Biophys. Acta <u>1090</u> :1-8
CX	Tominaga, 1989, "A putative protein of a growth specific cDNA from BALB/c-3T3 cells is highly similar to the extracellular portion of mouse interleukin 1 receptor", FEBS Lett. <u>258</u> :301-304
CY	Tsuyuki et al., 1997, "Costimulation through B7-2 (CD86) is required for the induction of a lung mucosal T helper cell 2 (Th2) immune response and altered airway responsiveness", J. Exp. Med. <u>185</u> :1671-1679
CZ	Volmer et al., 1994, "T lymphocytes derived from skin lesions of patients with psoriasis vulgaris express a novel cytokine pattern that is distinct from that of T helper type 2 cells", Eur. J. Immunol. <u>24</u> :2377-2382
DA	Wang et al., 1992, "IL-4 activates a distinct signal transduction cascade from IL-3 in factor-dependent myeloid cells", EMBO J. <u>11</u> :4899-4908
DB	Werenskiold, 1992, "Characterization of a secreted glycoprotein of the immunoglobulin superfamily inducible by mitogen and oncogene", Eur. J. Biochem. <u>204</u> :1041-1047
DC	Werenskiold et al., 1989, "Induction of a mitogen-responsive gene after expression of the Ha-ras oncogene in NIH 3T3 fibroblasts", Mol. Cell. Biol. <u>9</u> :5207-5214

DD	Wierenga et al., 1990, "Evidence for compartmentalization of functional subsets of CD4 <sup>+</sup> T lymphocytes in atopic patients", J. Immunol. <u>144</u> :4651-4656				
DE	Yamamura et al., 1991, "Defining protective responses to pathogens: Cytokine profiles in leprosy lesions", Science <u>254</u> :277-279				
DF	Yanagisawa et al., 1993, "Presence of a novel primary response gene ST2L, encoding a product highly similar to the interleukin 1 receptor type 1", FEBS Lett. <u>318</u> :83-87				
DG	Young et al., 1992, "Expression of cytolytic mediators by synovial fluid lymphocytes in rheumatoid arthritis", Am. J. Path. <u>140</u> :1261-1268				
DH	Zheng and Flavell, 1997, "The transcription factor GATA-3 is necessary and sufficient for Th2 cytokine gene expression in CD4 T cells", Cell <u>89</u> :587-596				
DI	GenBank Accession No. E07716, cDNA encoding human ST2, published June 1994				
<table border="1"> <tr> <td>EXAMINER</td> <td>DATE CONSIDERED</td> </tr> <tr> <td></td> <td>9-23-05</td> </tr> </table>		EXAMINER	DATE CONSIDERED		9-23-05
EXAMINER	DATE CONSIDERED				
	9-23-05				
<p>*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.</p>					